SEQUENCE LISTING

<110> Delansorne, Rémi
       Bonnet, Paule
       Paris, Jacques

<120> Pharmaceutical compositions based on alpha-cyclodextrin
       for the oral administration of LH-RH analogues

<130> H20058-5US

<140> 09/787,436
<141> 2000-03-17

<150> PCT/EP99/07389
<151> 1999-09-23

<150> EP98402403.4
<151> 1998-09-30

<160> 7

<170> PatentIn Ver. 2.1

<210> 1
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:LH-RH analogue

<220>
<221> SITE
<222> (1)
<223> Xaa is pGlu, D-pGlu, Sar, AcSar, Pro, Ser, D-Ser, Ac-D-Ser,
       Thr, D-Thr, Ac-D-Thr or an optionally substituted and/or acylated
       aromatic D-amino acid

<220>
<221> SITE
<222> (2)
<223> Xaa is His or an optionally substituted aromatic D-amino acid

<220>
<221> SITE
<222> (3)
<223> Xaa is an optionally substituted aromatic L- or D-amino acid

<220>
<221> SITE
<222> (4)
<223> Xaa is Ala, Ser, D-Ser, MeSer, Ser(OBut), Ser(OBzl) or Thr

<220>
<221> SITE
<222> (5)
<223> Xaa is an optionally substituted aromatic L-amino acid
       or an optionally substituted basic L- or D-amino acid

<220>

```
<221> SITE
<222> (6)
<223> Xaa is Gly, (S)-spirolactam-Pro, D-Pro, D-Ser, D-Thr,
      D-Cys, D-Met, D-Asn, D-Pen, D-(S-Me)Pen, D-(S-Et)Pen,
      D-Ser(OBut), D-Asp(OBut), D-Glu(OBut), D-Thr(OBut),
      D-Cys(OBut), D-Ser(OR1) where R1 is a sugar moiety

<220>
<221> SITE
<222> (6)
<223> Xaa is an aza-amino acid, D-His which may be substituted on the
      imidazole ring by a (C1-C6)alkyl, a (C2-C7)acyl or a
      benzyl group, an aliphatic D-amino acid with a (C1-C8)-
      alkyl or a (C3-C6)cycloalkyl side chain

<220>
<221> SITE
<222> (6)
<223> Xaa is an optionally substituted aromatic D-amino acid,
      D-cyclohexadienyl-Gly, D-perhydronaphthyl-Ala, D-perhydrodiphenyl-
Ala
      or an optionally substituted basic L- or D-amino acid

<220>
<221> SITE
<222> (7)
<223> Xaa is a linear, branched or cyclic aliphatic L-amino
      acid of 3 to 20 carbon atoms which may be N-alpha-
      substituted by a (C1-C4)alkyl group optionally substituted
      by one or several fluorine atoms

<220>
<221> SITE
<222> (8)
<223> Xaa is an optionally substituted basic L- or D-amino acid

<220>
<221> SITE
<222> (10)
<223> Xaa is GlyNH2, D-AlaNH2 or azaGlyNH2

<400> 1
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Pro Xaa
  1               5                   10

<210> 2
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:LH-RH analogue

<220>
<221> SITE
<222> (1)
<223> Xaa is pGlu, Sar or AcSar

<220>
<221> SITE
<222> (3)
```

<223> Xaa is an optionally substituted aromatic L-amino acid

<220>
<221> SITE
<222> (4)
<223> Xaa is Ala, Ser, D-Ser, MeSer, Ser(OBut), Ser(OBzl) or Thr

<220>
<221> SITE
<222> (5)
<223> Xaa is an optionally substituted aromatic L-amino acid

<220>
<221> SITE
<222> (6)
<223> Xaa is Gly, (S)-spirolactam-Pro, D-Pro, D-Ser, D-Thr,
      D-Cys, D-Met, D-Pen, D-(S-Me)Pen, D-(S-Et)Pen,
      D-Ser(OBut), D-Asp(OBut), D-Glu(OBut), D-Thr(OBut),
      D-Cys(OBut), D-Ser(OR1) where R1 is a sugar moiety

<220>
<221> SITE
<222> (6)
<223> Xaa is an aza-amino acid, D-His which may be substituted on the
      imidazole ring by a (C1-C6)alkyl or a benzyl group, an aliphatic
      D-amino acid with a (C1-C8)alkyl or a (C3-C6)cycloalkyl side
chain,
      an optionally substituted aromatic D-amino acid

<220>
<221> SITE
<222> (6)
<223> Xaa is D-cyclohexadienyl-Gly, D-perhydronaphthyl-Ala,
      D-perhydrodiphenyl-Ala or an optionally substituted basic D-amino
acid

<220>
<221> SITE
<222> (7)
<223> Xaa is a linear, branched or cyclic aliphatic L-amino
      acid of 3 to 20 carbon atoms which may be N-alpha-
      substituted by a (C1-C4)alkyl group optionally substituted
      by one or several fluorine atoms

<220>
<221> SITE
<222> (8)
<223> Xaa is an optionally substituted basic L-amino acid

<220>
<221> SITE
<222> (10)
<223> Xaa is GlyNH2 or azaGlyNH2

<400> 2
Xaa His Xaa Xaa Xaa Xaa Xaa Xaa Pro Xaa
  1   5   10

<210> 3
<211> 10
<212> PRT

<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:LH-RH analogue

<220>
<221> SITE
<222> (1)
<223> Xaa is pGlu

<220>
<221> SITE
<222> (3)
<223> Xaa is as defined for SEQ ID NO:2

<220>
<221> SITE
<222> (5)
<223> Xaa is as defined for SED ID NO:2

<220>
<221> SITE
<222> (6)
<223> Xaa is as defined for SED ID NO:2

<220>
<221> SITE
<222> (7)
<223> Xaa is Leu, Tle, Nle, Hol, Npg, Cha or Ada, which may
      be N-alpha-substituted by a methyl or ethyl group
      optionally substituted by one or several fluorine atoms

<220>
<221> SITE
<222> (10)
<223> Xaa is as defined for SEQ ID NO:2

<400> 3
Xaa His Xaa Ser Xaa Xaa Xaa Arg Pro Xaa
  1   5   10

<210> 4
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:LH-RH analogue

<220>
<221> SITE
<222> (1)
<223> Xaa is pGlu

<220>
<221> SITE
<222> (3)
<223> Xaa is Phe, Tyr, Trp, 2MeTrp, HPhe, HTyr, Nal, 1Nal,
      Bal, Pal, 4Pal or pClPhe

<220>

```
<221> SITE
<222> (5)
<223> Xaa is Phe, Tyr, Trp, 2MeTrp, HPhe, HTyr, Nal, 1Nal,
      Bal, Pal, 4Pal or pClPhe

<220>
<221> SITE
<222> (6)
<223> Xaa is (S)-spirolactam-Pro, Gly, D-Pro, D-Ser(OBut),
      D-Asp(OBut), D-Glu(OBut), D-Thr(OBut), D-Cys(OBut),
      D-His, D-His(Bzl), D-Ala, D-Leu, D-Tle, D-Nle, D-Hol,
      D-Npg, D-Cha, D-Phe, D-HPhe, D-Tyr, D-HTyr, D-Trp

<220>
<221> SITE
<222> (6)
<223> Xaa is D-2MeTrp, D-Nal, D-1Nal, D-Bal, D-Pal, D-4Pal, D-pClPhe
      D-cyclohexadienyl-Gly, D-perhydronaphtyl-Ala,
      D-perhydrodiphenyl-Ala or D-APhe optionally substituted
      by an aminotriazolyl group

<220>
<221> SITE
<222> (7)
<223> Xaa is Leu, Npg or Cha, which may be N-alpha-substituted
      by a methyl group

<220>
<221> SITE
<222> (10)
<223> Xaa is GlyNH2 or azaGlyNH2

<400> 4
Xaa His Xaa Ser Xaa Xaa Xaa Arg Pro Xaa
 1               5                   10

<210> 5
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:LH-RH analogue

<220>
<221> SITE
<222> (1)
<223> Xaa is pGlu

<220>
<221> SITE
<222> (6)
<223> Xaa is (S)-spirolactam-Pro, D-Leu, D-Ala, D-Nal,
      D-Phe, D-Ser(OBut) or D-Trp

<220>
<221> SITE
<222> (7)
<223> Xaa is Leu, MeLeu, Npg or MeNpg

<220>
```

```
<221> SITE
<222> (10)
<223> Xaa is GlyNH2, azaGlyNH2 or -NC2H5

<400> 5
Xaa His Trp Ser Tyr Xaa Xaa Arg Pro Xaa
 1               5                   10

<210> 6
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:LH-RH analogue


<220>
<221> SITE
<222> (1)
<223> Xaa is pGlu, D-pGlu, Sar, AcSar, Pro thereof, Ser, D-Ser,
      Ac-D-Ser, Thr, D-Thr, Ac-D-Thr or an optionally substituted
      and/or acylated aromatic D-amino acid

<220>
<221> SITE
<222> (2)
<223> Xaa is an optionally substituted aromatic D-amino acid

<220>
<221> SITE
<222> (3)
<223> Xaa is an optionally substituted aromatic L- or D-amino acid

<220>
<221> SITE
<222> (4)
<223> Xaa is Ala, Ser, D-Ser, MeSer, Ser(OBut), Ser(OBzl) or Thr

<220>
<221> SITE
<222> (5)
<223> Xaa is an optionally substituted aromatic L-amino acid
      or an optionally substituted basic L- or D-amino acid

<220>
<221> SITE
<222> (6)
<223> Xaa is Gly, (S)-spirolactam-Pro, D-Pro, D-Ser, D-Thr,
      D-Cys, D-Met, D-Asn, D-Pen, D-(S-Me)Pen, D-(S-Et)Pen,
      D-Ser(OBut), D-Asp(OBut), D-Glu(O-But), D-Thr(O-But),
      D-Cys(O-But), D-Ser(O-R1) where R1 is a sugar moiety

<220>
<221> SITE
<222> (6)
<223> Xaa is an aliphatic D-amino acid with a (C1-C8)alkyl or a(C3-C6)
      cycloalkyl side chain, an optionally substituted aromatic D-amino
acid
      D-cyclohexadienyl-Gly, D-perhydronaphthyl-Ala, D-perhydrodiphenyl-
Ala
```

or an optionally substituted basic L- or D-amino acid

<220>
<221> SITE
<222> (7)
<223> Xaa is a linear, branched or cyclic aliphatic L-amino acid of 3 to
       20 carbon atoms which may be N-alpha-substituted by a (C1-C4)alkyl
       group optionally substituted by one or several fluorine atoms

<220>
<221> SITE
<222> (8)
<223> Xaa is an optionally substituted basic L- or D-amino acid

<220>
<221> SITE
<222> (10)
<223> Xaa is GlyNH2 or D-AlaNH2

<400> 6
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Pro Xaa
  1   5   10

<210> 7
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:LH-RH analogue

<220>
<221> SITE
<222> (1)
<223> Xaa is Ac-D-Nal

<220>
<221> SITE
<222> (2)
<223> Xaa is D-pClPhe

<220>
<221> SITE
<222> (3)
<223> Xaa is D-Pal

<220>
<221> SITE
<222> (5)
<223> Xaa is Tyr, HTyr, MeTyr, MeHTyr, NicLys or IprLys

<220>
<221> SITE
<222> (6)
<223> Xaa is (S)-spirolactam-Pro, D-Arg, D-NicLys, D-IprLys,
       D-Cit, D-HCit or D-Asn

<220>
<221> SITE
<222> (7)

```
<223> Xaa is Leu, MeLeu, Npg or MeNpg

<220>
<221> SITE
<222> (8)
<223> Xaa is Arg, NicLys or IprLys

<220>
<221> SITE
<222> (10)
<223> Xaa is DAlaNH2

<400> 7
Xaa Xaa Xaa Ser Xaa Xaa Xaa Xaa Pro Xaa
  1               5                   10
```